

—Review—

## ***In Vitro Culture and Evaluation of Embryos for Production of High Quality Bovine Embryos***

Hiroyuki Abe<sup>1\*</sup>, Hitoshi Shiku<sup>2</sup>, Shigeo Aoyagi<sup>3</sup> and Hiroyoshi Hoshi<sup>4</sup>

<sup>1</sup>Tohoku University Biomedical Engineering Research Organization, 2-1 Seiryō, Aoba-ku, Sendai, Miyagi 980-8575, Japan

<sup>2</sup>Department of Biomolecular Engineering, Graduate School of Engineering, Tohoku University, Sendai, Miyagi 980-8579, Japan

<sup>3</sup>Research & Development Department, Hokuto Denko Corporation, Atsugi, Kanagawa 243-0801, Japan

<sup>4</sup>Research Institute for the Functional Peptides, Yamagata 990-0823, Japan

Recent advances in technologies associated with assisted reproduction in domestic animals have been remarkable. In cattle, immature oocytes can be developed to the blastocyst stage by *in vitro*-production (IVP) techniques in various formulations of media and different culture conditions [1–5], but many technical problems still remain. For example, the cryotolerance and rates of pregnancy of embryos produced by IVP techniques were not high [6, 7]. Comparative studies demonstrated that the quality of bovine embryos produced by IVP systems was lower than that of *in vivo*-derived embryos [8, 9]. Embryo quality is an important determinant of the success of practical embryo transfer. Accurate evaluation of embryo quality prior to the embryo transfer contributes to improvement in rates of pregnancy. Therefore, successful assisted reproduction in animals requires culture techniques for production of high quality embryos and an accurate method for assessment of embryo quality. This review introduces *in vitro* culture systems for production of high quality bovine embryos and a novel method, with scanning electrochemical microscopy, to assess the embryo quality.

### **Culture Systems for Embryo Production**

Bovine immature oocytes can be developed to the blastocyst stage by an IVP technique. Bovine embryos are routinely cultured in serum-supplemented media with or without somatic cell co-cultures, because serum contains beneficial substances for embryonic

development such as growth factors, hormones, anti-oxidative compounds, and chelators of heavy metals [10]. But the biological activities of sera vary from lot to lot and there are potential risks of virus or mycoplasma infections. Furthermore, serum has a biphasic effect on bovine embryo development, inhibiting the first cleavage division but stimulating blastocyst development [1, 11]. Some studies have shown that the addition of serum to culture medium can cause morphological abnormalities in embryos [10, 12–14] and affect gene expression in bovine matured oocytes [15] and in pre-implantation embryos [16]. Serum is also suspected of contributing to the large offspring syndrome in domestic animal species [13, 17, 18]. Obviously, serum is a key factor in the culture environment affecting embryo quality, but because of the negative effects of serum, culture systems with serum-free medium, which is capable of efficient production of high quality bovine embryos, are desirable.

In order to formulate a serum-free medium for bovine embryo production, we conducted investigations to determine the factors required for oocyte maturation and subsequent embryonic development. Transforming growth factor- $\alpha$  (TGF- $\alpha$ ) and epidermal growth factor (EGF), as well as luteinizing hormone (LH) and follicle stimulating hormone (FSH) have been shown to enhance cumulus expansion and oocyte fertility [19]. Our study also showed that TGF- $\alpha$  and/or EGF are beneficial for obtaining matured oocytes of high quality for IVF. It is known that certain factors secreted by

somatic cells, such as oviduct epithelial cells and granulosa cells, are involved in embryonic development *in vitro* [20–22]. We investigated the embryotrophic factors in bovine granulosa cell-conditioned medium (BGC-CM) by means of a serum-free culture system. The embryotrophic activity of BGC-CM has been purified and identified as a tissue inhibitor of metalloproteinase-1 (TIMP-1) [23]. TIMP-1 enhanced the developmental rates from the morula to blastocyst stage and increased the numbers of cells in embryos at the blastocyst stage. Furthermore, it has been shown that both basic fibroblast growth factor (bFGF) and transforming growth factor- $\beta_1$  (TGF- $\beta_1$ ) increased the rate of bovine embryos developed to the blastocyst stage in a serum-free medium without somatic cell co-culture [24]. It has also been shown that the combination of a low oxygen atmosphere (5% O<sub>2</sub>) and low glucose medium (2.22 mM) improved the development of 1–2 cell embryos to blastocysts in a defined medium without somatic cell co-cultures [25].


On the basis of the abovementioned studies, two serum-free media (IVMD101 and IVD101) have been developed for bovine oocyte maturation and embryo culture [26]. The formulations of these serum-free media have been described in other papers [27, 28]. IVMD101 medium is used for both *in vitro* oocyte maturation and embryo culture in the presence of cumulus/granulosa cells under an atmosphere of 5% CO<sub>2</sub> in air. IVD101 medium is used for embryo culture in the absence of somatic cell co-culture under a low oxygen condition (5% O<sub>2</sub>).

### Physiological Evaluations of Bovine Embryos

These serum-free media showed improvement in the developmental rates and quality of bovine blastocysts produced by IVP systems [26]. The proportion of embryos developing to the blastocyst stage in IVMD101 medium with bovine cumulus/granulosa cells (BCGC) co-cultures (36.5%) and IVD101 medium without BCGC co-cultures under a low oxygen condition (37.1%) was significantly higher than in serum-containing medium (TCM-199 + 5% calf serum; CS) with BCGC co-cultures (25.1%). The mean numbers of cells in Day 7 blastocysts developed in IVMD101 (179.5 cells) and IVD101 (177.1 cells) media were greater than that in the serum-supplemented medium (145.7 cells). The survival rates of blastocysts produced in IVMD101 medium (73.3%) and IVD101 medium (60.0%) based on hatching after 72 h of post-thaw culture were superior to that of blastocysts produced in TCM-199 + CS medium

(48.1%).

It is well known that embryo transfer of *in vitro*-derived bovine embryos often results in the production of bovine calves with abnormalities (large offspring syndrome), such as heavier than normal birth weight and a high incidence of dystocia [7, 29–31]. Bovine embryos exposed to a variety of unusual environments prior to the blastocyst have resulted in the development of unusually large offspring. The large offspring syndrome has been started in bovine and ovine embryos after exposure to unusual environments both *in vivo* and *in vitro*. Four different situations have been identified that result in the syndrome: *in vitro* embryo culture, asynchronous embryo transfer into an advanced uterine environment, nuclear transfer, and maternal exposure to an excessively high urea diet [29–32]. Neither the environmental factors inducing the large offspring syndrome nor the mechanisms of perturbation occurring in the early embryo and manifesting themselves in the fetus have been identified. Although the nature of the perturbing environmental factors has not yet been identified, serum-supplemented culture conditions are suspected of contributing to the large offspring syndrome in sheep [13, 17, 18]. Many components of serum such as growth factors, free radicals, ammonia, and progesterone may be candidate factors perturbing environments [33]. Therefore, the normality of calves derived from embryos produced in serum-free media is of great interest. Pregnancy, parturition and calf mortality rates of Japanese Black calves were compared after transfer of embryos developed in either serum-free media (IVMD101 and IVD101) or serum-supplemented media (TCM-199 and HPM199 + 5% CS) [28]. The calving rate of recipients receiving embryos from serum-free cultures (39.6%) was slightly heavier than those receiving embryos from serum-supplemented cultures (32.8%). Although the calving rate and abortion rate for embryos in serum-free and serum-supplemented cultures were not very different (85.2% vs 86.4%; 14.8% vs 13.6%, respectively), calf mortality for embryos from serum-supplemented cultures was greater than that for embryos from serum-free cultures (13.6% vs 4.9%). Average birth weights of calves for embryos from serum-supplemented cultures were slightly higher than those for embryos from serum-free cultures (male: 37.5 kg vs 34.0 kg; female: 36.1 kg vs 32.1 kg, respectively). Although these mean birth weights of calves were within the previously reported normal range of birth weight for Japanese Black calves produced from *in vitro* produced embryos [37], the birth weights of calves from embryos derived in serum-



**Fig. 1.** Electron micrographs of bovine morulae cultured in serum-supplemented medium (TCM-199 + CS; a, c) and serum-free medium (IVMD101; b, d). a: Note cells filled with numerous lipid droplets (LD) of high electron density. b: Lipid droplets were present in cytoplasm, but were not abundant. c: Note immature mitochondria (M) showing ovoid structures with a hooded appearance and a dark electron-dense matrix. d: Note presence of elongated mitochondria with transverse cristae. N, nucleus; V, vesicles; ZP, zona pellucida. Bars: a, b = 2  $\mu$ m, c, d = 1  $\mu$ m.

supplemented culture (18–74 kg) were more variable than those from embryos developed in serum-free cultures (23–45 kg).

The serum-free media described in this paper are useful for the production of high quality bovine embryos for transfer. Comparative studies of bovine embryos cultured in different culture systems with serum-free or serum-containing media are examined in more detail below.

### Morphological Evaluations of Bovine Embryos

Conspicuous differences in the morphological features of bovine embryos cultured in serum-free and serum-supplemented media have been observed [14]. Morulae and blastocysts cultured in serum-supplemented medium (TCM-199 + 5% CS) were darker and contained numerous cytoplasmic inclusions compared to those cultured in serum-free media (IVMD101 and IVD101). Histochemical examinations revealed that bovine embryos cultured in serum-containing medium had more, much larger lipid droplets in the cytoplasm than those cultured in serum-free media. Cytoplasmic lipid droplets increased substantially in embryos from the morula to the

blastocyst stages when cultured in serum-containing medium [38].

Electron microscopic examinations showed major differences in the ultrastructural features of the morulae and blastocysts from serum-free and serum-supplemented cultures [14]. The blastomeres of morulae and blastocysts grown in serum-supplemented medium contained many large, electron-dense lipid droplets, whereas those cultured in serum-free media had fewer (Fig. 1a, b). Morulae and blastocysts derived *in vivo* contained less cytoplasmic lipid, similar to those developed in serum-free cultures. These findings strongly suggest that the presence of serum in the culture medium may be the cause of the abnormal accumulation of cytoplasmic lipids in bovine embryos. Similar findings showed that ovine embryos cultured in human serum-supplemented medium had abundant cytoplasmic lipid droplets [10, 13].

It is interesting to note that a conspicuous difference in the maturation of mitochondria has been observed between embryos cultured in serum-free and serum-supplemented media [14, 38]. There were numerous elongated mitochondria in the morulae developed in serum-free media, whereas many of the mitochondria of the morulae cultured in serum-containing medium were

spherical or ovoid, indicative of an immature form (Fig. 1c, d). Changes in mitochondrial ultrastructure and development were observed in bovine embryos from morula to early blastocyst stages in other studies [39–42]. It seems that the presence of serum had adverse effects on the structure and development of mitochondria. Similar mitochondrial changes were found in ovine embryos cultured in synthetic oviduct fluid (SOF) medium supplemented with human serum [43].

### Cryotolerance and Lipid Accumulation

It is of great interest to clarify the correlation between the lipid accumulation and cryotolerance. Comparative studies of *in vivo* and *in vitro* produced embryos showed substantial differences in their morphology, metabolism, and overall embryo quality as assessed by the numbers of cells and frozen embryo viability [3, 8, 9, 38]. Although bovine blastocysts produced by an IVP technique have yielded live calves after transfer [44, 45], standard cryopreservation methods which yield high survival rates for *in vivo*-derived bovine embryos, usually result in low survival rates for *in vitro*-developed embryos, suggesting that *in vitro*-produced bovine embryos are more sensitive to cryopreservation than *in vivo*-derived embryos [6, 46, 47]. The displacement of intracellular lipids by centrifugation significantly improved the survival of *in vitro* developed embryos, suggesting that those lipids were partially responsible for the sensitivity to chilling and freezing [48, 49]. To evaluate the correlation between the lipid accumulation in bovine embryos and the cryotolerance of embryos, we examined the tolerance of cryopreservation by embryos at different developmental stages grown in either serum-free or serum-containing media [38]. The post-thaw viability and hatching rates of bovine embryos developed in serum-free media from the morula to the blastocyst stages was significantly higher than those developed in serum-containing medium, suggesting that bovine embryos with high lipid content are more sensitive to cryopreservation procedures. In summary, sensitivity of bovine embryos produced by an IVP technique to cryopreservation could be improved by using serum-free cultures. Therefore, serum-free media are beneficial for the production of high quality, cryogenically preserved bovine embryos.

The mechanism of cytoplasmic lipid accumulation in bovine embryos cultured in serum-containing medium is still unclear. It is well known that cells in cultures can readily take up fatty acids, phospholipids, and

triglycerides from serum-supplemented medium and most of the lipids are derived from triglycerides contained in serum lipoproteins [50]. Bovine embryos cultured in a serum-supplemented medium showed a different fatty acid composition to those cultured in a serum-free medium [51]. The fatty acid composition of embryos grown in calf serum-containing medium was similar to those of calf serum, suggesting that the accumulation of long-chain saturated and unsaturated fatty acids in serum-exposed embryos is from the serum itself. It has been reported that excessive accumulation of triglycerides occurs in bovine embryos cultured in serum-containing medium [52]. Furthermore, a recent study by us showed that bovine embryos developed in serum-free media supplemented with bovine serum plasma lipoproteins contained numerous lipid droplets in the cytoplasm (unpublished results). These findings suggest that the source of cytoplasmic lipids in bovine embryos cultured in serum-supplemented medium is in the lipoproteins of the serum itself.

### Novel Technique for Quality Evaluation of Embryos

The quality of embryos prior to embryo transfer affects the pregnancy rates. Several methods for embryo evaluation have been developed. Dye exclusion tests [53], measurement of enzyme activity [54], glucose uptake [55], and live-dead stains [56] are useful in predicting embryo survival after transfer, but most of these methods are harmful and therefore not useful for the evaluation of individual embryos. In the past, morphological evaluation has been widely used to evaluate embryo quality because it is non-invasive and useful in predicting pregnancy rates of groups of bovine embryos after transfer [57–61]. Based on morphological evaluations, embryos of good quality yielded high pregnancy rates and poor embryos resulted in low success rates. Nevertheless, in some cases, embryos of low quality produced a pregnancy, whereas many good embryos as judged by morphological observation failed to result in pregnancy. Furthermore, morphological evaluations remain one of the most subjective and quantitative aspects of embryo transfer because categorization standards vary among investigators.


A recent study by us showed conspicuous differences in the ultrastructural features of morulae of high and low quality [62]. The morulae classified as low quality (fair and poor) by morphological classification contained nucleoli with low transcriptional activity, a large number

of lipid droplets, and immature mitochondria, suggesting that low quality embryos have low metabolic activities, including oxygen consumption. In embryos, the maturation of mitochondria is associated with increases in metabolism reflected in their oxygen consumption [63] and CO<sub>2</sub> production [64], and it appears to be related to the depletion of stored products (cytoplasmic lipids) so that mitochondrial function (oxygen consumption) may be an important factor in embryo quality.

It is well known that metabolic processes drastically change during embryonic development, as indicated by genome activation and large increases in protein synthesis. Radioisotope labeling techniques have been widely used to quantify glucose, pyruvate, lactate, carbon dioxide and amino acids in metabolic pathways, and detection of metabolites originating in a single embryo have been attempted [64–67]. Oxygen consumption is an ubiquitous parameter also used to gain valuable information on metabolic mechanisms. Oxygen consumption of mammalian embryos has been studied with various methods such as Cartesian diver [63], spectrophotometrics [68–70], fluorescence [71–73], and electrochemical techniques [74–78].

Recently, we succeeded in non-invasively and quantitatively determining oxygen consumption of individual bovine embryos cultured in the serum-free medium (IVD101) by scanning electrochemical microscopy (SECM) [79]. SECM is a technique in which the tip of a microelectrode is used to scan and monitor the local distribution of electro-active species (O<sub>2</sub>) near the sample surface (embryo). With this technique, oxygen consumption of single, identical bovine embryos at different developmental stages has been monitored. The morulae with the higher oxygen consumption on day 6 developed into blastocysts of larger sizes and higher oxygen consumption on day 8 [79]. The morulae with higher oxygen consumption have a strong potential for further development into expanded and/or hatched blastocysts, thereby increasing the possibility that the SECM procedure could be used to non-invasively assess the embryo quality and the subsequent developmental potential of embryos.

Although SECM may be useful for judging bovine embryo quality, the SECM measuring procedure requires quite a bit of skill, such as gently holding a single embryo with a micropipette positioned with a micromanipulator and aligning the microelectrode very close to the embryo surface. Recently, we designed a new SECM measuring procedure which can be easily used by a non-electrochemist. This new SECM



**Fig. 2.** A new scanning electrochemical microscopy (SECM) system (a) and a plate for measuring respiratory activity of bovine embryos (b). a: SECM includes a measuring instrument on the inverted optical microscope stage (A), a potentiostat (B), notebook computer, and monitor. b: The plate has six cone-shaped microwells (arrow). Individual embryos are transferred into a microwell filled with culture medium. The embryo sinks down to the bottom of the well, remaining at the lowest point. The oxygen concentration profiles are measured according to the method by Shiku et al. (unpublished results).

measuring system includes a measuring instrument on an inverted optical microscope stage, a potentiostat (Hokuto Denko Co., Tokyo, Japan), and a notebook computer as controller and analyzer (Fig. 2a). The measuring instruments are covered with a plastic chamber and humidified 5% CO<sub>2</sub> and 95% air gas can be maintained. The motor driven XYZ-stage is located on the microscope stage and this stage and the potentiostat are controlled by a computer. The oxygen consumption rate of embryos is calculated by newly designed software. To easily handle many embryos in a short time, a plate with cone-shaped microwells was designed (Fig. 2b). This plate is designed to amplify oxygen concentration changes due to respiration by the embryo. Using this modified procedure, oxygen consumption has been monitored at various developmental stages of single, identical bovine embryos cultured in IVD101 medium (Table 1). Oxygen consumption rates (*F*) of the single embryos were low

**Table 1.** Oxygen consumption rates of the bovine embryos cultured in IVD101 medium at various stages of development

Embryonic stage	No. of embryos measured	Oxygen consumption rate ( $F \times 10^{14} / \text{mol s}^{-1}$ )
2 cell	15	$0.46 \pm 0.05^a$
4 cell	17	$0.45 \pm 0.03^a$
8 cell	18	$0.46 \pm 0.02^a$
Morula	48	$1.03 \pm 0.05^b$
Blastocyst	55	$1.86 \pm 0.07^c$

Values with different superscripts in each column differ significantly ( $P < 0.05$ ).

**Table 2.** Relationship between the oxygen consumption and the further developmental potential of the bovine morulae cultured in IVD101 medium

Oxygen consumption rate ( $F \times 10^{14} / \text{mol s}^{-1}$ )	No. of embryos* measured	No. of embryos developed to blastocyst after 96 h (%)	No. of blastocysts expanded or hatched after 96 h (%)
$F \geq 1.0$	56	50 (89.3)	35 (62.5)
$0.8 \leq F < 1.0$	44	34 (77.3)	20 (50.0)
$F < 0.8$	107	49 (45.8)	28 (26.2)

\*: Bovine embryos at the morula stage were selected at day 6 after in vitro fertilization. After measurement by a modified scanning electrochemical microscopy, the embryos were incubated in IVD101 medium for 4 days in a humidified atmosphere of 5% CO<sub>2</sub>/5% O<sub>2</sub>/90% N<sub>2</sub> at 38.5°C.

from 2-cell to 8-cell stages ( $0.48\text{--}0.55 \times 10^{-14} \text{ mol s}^{-1}$ ). An increase in the oxygen consumption rate was found at the morula ( $1.03 \times 10^{-14} \text{ mol s}^{-1}$ ) stage and blastocysts showed an even higher oxygen consumption rate ( $1.86 \times 10^{-14} \text{ mol s}^{-1}$ ). The morulae with the highest oxygen consumption have the strongest potential for further development into expanded and/or hatched blastocysts (Table 2). Furthermore, we found that the oxygen consumption rates of bovine embryos (morulae and blastocysts) produced in serum-free medium (IVD101) were higher than those of embryos cultured in serum-supplemented medium (HPM199 + CS), indicating a correlation between respiratory activity and development of mitochondria (unpublished results). These results demonstrate that the novel SECM procedures are very useful for non-invasively and easily evaluating the oxygen consumption of bovine embryos. This SECM system may have a wide application for judging embryo quality and may be an ideal method for noninvasive quality evaluation of embryos in the future.

### Conclusions

With serum-free culture for IVF and IVM bovine embryos can offer several advantages over the cultures with serum-supplemented media. The improved serum-free media (IVMD101 and IVD101) are especially useful

for the production of higher quality bovine embryos for cryopreservation. Furthermore, it has been demonstrated that there is a correlation between oxygen consumption and embryo quality. Scanning electrochemical microscopy can be non-invasively applied to measure oxygen consumption of single bovine embryos. This novel method may be useful to assess the quality of bovine embryos and their developmental potential. The combination of novel culture systems and evaluation techniques for bovine embryos may contribute to improvements in the rates of pregnancy with bovine embryos produced by IVP technique.

### Acknowledgments

The authors thank T. Satoh, K. Kobayashi, M. Kacchi, and S. Yamashita for assisting with the development of serum-free media and Y. Ustumi, M. Matsudaira, T. Shiraishi, and Prof. T. Matsue for the design of the new SECM system. This research was supported by the Research Fund of the Bio-Oriented Technology Research Advancement Institution (BRAIN) of Japan, Yamagata Prefecture Joint/Research Project for Regional Intensive, Japan Science and Technology Corporation, and Japan Livestock Technology Association.

## References

- 1) Pinyopummintr, T. and Bavister, B.D. (1991): In vitro-matured/in vitro-fertilized bovine oocytes can develop into morulae/blastocysts in chemically defined protein-free culture media. *Biol. Reprod.*, 45, 736–742.
- 2) Voelkel, S.A. and Hu, Y.X. (1992): Effect of gas atmosphere on the development of one-cell bovine embryos in two culture systems. *Theriogenology*, 37, 1117–1131.
- 3) Shamsuddin, M., Larsson, B., Gustafsson, H. and Rodriguez-Martinez, H. (1993): In vitro development up to hatching of bovine in vitro-matured and fertilized oocytes with or without support from somatic cells. *Theriogenology*, 39, 1067–1079.
- 4) Shamsuddin, M. and Larsson, B. (1993): In vitro development of bovine embryos after fertilization using semen from different donors. *Reprod. Dom. Anim.*, 28, 77–84.
- 5) Shamsuddin, M., Larsson, B. and Rodriguez-Martinez, H. (1993): Culture of bovine IVM/IVF embryos up to blastocyst stage in defined medium using insulin, transferrin and selenium or growth factors. *Reprod. Dom. Anim.*, 28, 209–210.
- 6) Leibo, S.P. and Loskutoff, N.M. (1993): Cryobiology of in vitro-derived bovine embryos. *Theriogenology*, 39, 81–94.
- 7) Schmidt, M., Greve, T., Avery, B., Beckers, J.F., Sulon, J. and Hansen, H.B. (1996): Pregnancies, calves and calf viability after transfer of in vitro produced bovine embryos. *Theriogenology*, 46, 527–539.
- 8) Greve, T., Avery, B. and Callesen, H. (1993): Viability of in-vivo and in-vitro produced bovine embryos. *Reprod. Dom. Anim.*, 28, 164–169.
- 9) Massip, A., Mermillod, P. and Dinnyes, A. (1995): Morphology and biochemistry of in-vitro produced bovine embryos: implications for their cryopreservation. *Hum. Reprod.*, 10, 3004–3011.
- 10) Gardner, D.K. (1994): Mammalian embryo culture in the absence of serum or somatic cell support. *Cell Biol. Int.*, 18, 1163–1178.
- 11) Van Langendonck, A., Donnay, I., Schuurbiers, N., Auquier, P., Carolan, C., Massip, A. and Dessy, F. (1997): Effects of supplementation with fetal calf serum on development of bovine embryos in synthetic oviduct fluid medium. *J. Reprod. Fertil.*, 109, 87–93.
- 12) Gardner, D.K., Lane, M., Spitzer, A. and Batt, P.A. (1994): Enhanced rates of cleavage and development for sheep zygotes cultured to the blastocyst stage in vitro in the absence of serum and somatic cells: amino acids, vitamins, and culturing embryos in groups stimulate development. *Biol. Reprod.*, 50, 390–400.
- 13) Thompson, J.G., Gardner, D.K., Pugh, P.A., McMillan, W.H. and Tervit, H.R. (1995): Lamb birth weight is affected by culture system utilized during in vitro pre-elongation development of ovine embryos. *Biol. Reprod.*, 53, 1385–1391.
- 14) Abe, H., Yamashita, S., Itoh, T., Satoh, T. and Hoshi, H. (1999): Ultrastructure of bovine embryos developed from in vitro-matured and -fertilized oocytes: comparative morphological evaluation of embryos cultured either in serum-free or in serum-supplemented medium. *Mol. Reprod. Dev.*, 53, 325–335.
- 15) Rzucidlo, S.J., Gibbons, J. and Stice, S.L. (2001): Comparison by restriction fragment differential display RT-PCR of gene expression pattern in bovine oocytes matured in the presence or absence of fetal calf serum. *Mol. Reprod. Dev.*, 59, 90–96.
- 16) Wrenzycki, C., Herrmann, D., Carnwath, J.W. and Niemann, H. (1999): Alterations in the relative abundance of gene transcripts in preimplantation bovine embryos cultured in medium supplemented with either serum or PVA. *Mol. Reprod. Dev.*, 53, 8–18.
- 17) Holm, P., Walker, S.K. and Seamark, R.F. (1996): Embryo viability, duration of gestation and birth weight in sheep after transfer of in vitro matured and in vitro fertilized zygotes cultured in vitro or in vivo. *J. Reprod. Fertil.*, 107, 175–181.
- 18) Sinclair, K.D., McKvoy, T.G., Carolan, C., Maxfield, E.K., Maltin, C.A., Young, L.E., Wilmut, I., Robinson, J.J. and Broadbent, P.J. (1998): Conceptus growth and development following in vitro culture of ovine embryos in media supplemented with bovine sera. *Theriogenology*, 49, 218.
- 19) Kobayashi, K., Takagi, Y., Satoh, T., Hoshi, H. and Oikawa, T. (1992): Development of early bovine embryos to the blastocyst stage in serum-free conditioned medium from granulosa cells. *In Vitro Cell. Dev. Biol.*, 28A, 255–259.
- 20) Bavister, B.D. (1995): Culture of pre-implantation embryos: facts and artifacts. *Hum. Reprod. Update*, 1, 91–148.
- 21) Eyestone, W.H. and First, N.L. (1989): Co-culture of early cattle embryos to the blastocyst stage with oviducal tissue or in conditioned medium. *J. Reprod. Fertil.*, 85, 715–720.
- 22) Gandolfi, F. (1994): Autocrine, paracrine and environmental factors influencing embryonic development from zygote to blastocyst. *Theriogenology*, 41, 95–100.
- 23) Satoh, T., Kobayashi, K., Yamashita, S., Kikuchi, M., Sendai, Y. and Hoshi, H. (1994): Tissue inhibitor of metalloproteinases (TIMP-1) produced by granulosa and oviduct cells enhances in vitro development of bovine embryo. *Biol. Reprod.*, 50, 835–844.
- 24) Hoshi, H. (1996): In vitro production of bovine pre-implantation embryos in a serum-free medium. In: *Gene Expression and Nutrition in Animals: From Cells to Whole Body*. (Muramatsu, T. ed.), pp. 59–70, Research Signpost, Trivandrum, India.
- 25) Kobayashi, K., Yamashita, S. and Hoshi, H. (1994): Influence of epidermal growth factor and transforming growth factor- $\alpha$  on in vitro maturation of cumulus cell-enclosed bovine oocytes in a defined medium. *J. Reprod. Fertil.*, 100, 439–446.
- 26) Yamashita, S., Abe, H., Itoh, T., Satoh, T. and Hoshi, H. (1999): A serum-free culture system for efficient in vitro production of bovine blastocysts and their improved

- viability after freezing and thawing. *Cytotechnology*, 31, 1–9.
- 27) Abe, H. and Hoshi, H. (2003): Evaluation of bovine embryos produced in high performance serum-free media. *J. Reprod. Dev.*, 49, 193–202.
  - 28) Hoshi, H. (2003): In vitro production of bovine embryos and the application for embryo transfer. *Theriogenology*, 59, 675–685.
  - 29) Young, L.E., Sinclair, K.D. and Wilmot, I. (1998): Large offspring syndrome in cattle and sheep. *Rev. Reprod.*, 3, 155–163.
  - 30) Young, L.E., Butterwith, S.C. and Wilmot, I. (1996): Increased ovine fetal weight following transient asynchronous embryo transfer is not associated with increased placental weight at day 21 of gestation. *Theriogenology*, 45, 231.
  - 31) McEvoy, T.G., Robinson, J.J., Findley, P.A., Aitken, R.P. and Robertson, I.S. (1997): Dietary excesses of urea influence the viability and metabolism of preimplantation sheep embryos and may affect fetal growth among the survivors. *Anim. Reprod. Sci.*, 47, 71–90.
  - 32) Wilmot, I., Campbell, K.H.S. and Young, L.E. (1997): Modern reproduction technologies and transgenics. In: *The Genetics of Sheep*. (Piper, L. and Ruvinsky, A. eds.), pp. 395–411, CAB International, Oxford.
  - 33) Kaye, P.L. (1997): Preimplantation growth factor physiology. *Rev. Reprod.*, 2, 121–127.
  - 34) Farin, P. and Farin, C. (1997): Perspectives on large calves following transfer of embryos produced in vitro. *Embryo Transfer Newslett.*, 15, 15–19.
  - 35) Behboodi, E., Anderson, G.B., BonDurant, R.H., Cargill, S.L., Kreuzer, B.R., Medrano, J.F. and Murray, J.D. (1995): Birth of large calves that developed from in vitro-derived bovine embryos. *Theriogenology*, 44, 227–232.
  - 36) Kruij, T.A.M. and den Daas, J.H.G. (1997): In vitro produced and cloned embryos: effects on pregnancy, parturition and offspring. *Theriogenology*, 47, 43–52.
  - 37) Yazawa, S., Aoyagi, Y., Konishi, M. and Takedomi, T. (1997): Characterization and cytogenetic analysis of Japanese Black calves produced by nuclear transfer. *Theriogenology*, 48, 641–650.
  - 38) Abe, H., Yamashita, S., Satoh, T. and Hoshi, H. (2002): Accumulation of cytoplasmic lipid droplets in bovine embryos and cryotolerance of embryos developed in different culture systems using serum-free or serum-containing media. *Mol. Reprod. Dev.*, 61, 57–66.
  - 39) Abe, H., Otoi, T., Tachikawa, S., Yamashita, S., Satoh, T. and Hoshi, H. (1999): Fine structure of bovine morulae and blastocysts in vivo and in vitro. *Anat. Embryol.*, 199, 519–527.
  - 40) Brackett, B.G., Oh, Y.K., Evans, J.F. and Donawick, W.J. (1980): Fertilization and early development of cow ova. *Biol. Reprod.*, 23, 189–205.
  - 41) Mohr, L.R. and Trounson, A.O. (1981): Structural changes associated with freezing of bovine embryos. *Biol. Reprod.*, 25, 1009–1025.
  - 42) Plante, L. and King, W.A. (1994): Light and electron microscopic analysis of bovine embryos derived by in vitro and in vivo fertilization. *J. Assist. Reprod. Genet.*, 11, 515–529.
  - 43) Dorland, M., Gardner, D.K. and Trounson, A.O. (1994): Serum in synthetic oviduct fluid causes mitochondrial degeneration in ovine embryos. *J. Reprod. Fertil. (Abstract Series)*, 13, 25.
  - 44) Fukuda, Y., Ichikawa, M., Naito, K. and Toyoda, Y. (1990): Birth of normal calves resulting from bovine oocytes matured, fertilized, and cultured with cumulus cells in vitro up to the blastocyst stage. *Biol. Reprod.*, 42, 114–119.
  - 45) Zhang, L., Barry, D.M., Denniston, R.S., Bunch, T.D. and Godke, R.A. (1993): Birth of live calves after transfer of frozen-thawed bovine embryos fertilized in vitro. *Vet. Rec.*, 132, 247–279.
  - 46) Pollard, J.W. and Leibo, S.P. (1993): Comparative cryobiology of in vitro and in vivo derived bovine embryos. *Theriogenology*, 39, 287.
  - 47) Hasler, J.F., Henderson, W.B., Hurtgen, P.J., Jin, Z.Q., McCauley, A.D., Mower, S.A., Neely, B., Shuey, L.S., Stokes, J.E. and Trimmer, S.A. (1995): Production, freezing and transfer of bovine IVF embryos and subsequent calving results. *Theriogenology*, 43, 141–152.
  - 48) Leibo, S.P., Pollard, J.W. and Martino, A. (1995): Chilling and freezing sensitivity of “reassembled” in vitro-derived bovine embryos. *Theriogenology*, 43, 265.
  - 49) Ushijima, H., Yamakawa, H. and Nagashima, H. (1999): Cryopreservation of bovine pre-morula-stage in vitro matured/in vitro fertilized embryos after delipidation and before use in nucleus transfer. *Biol. Reprod.*, 60, 535–539.
  - 50) Spector, A.A. (1973): Fatty acid, glyceride, and phospholipid metabolism. In: *Growth, Nutrition and Metabolism of Cells in Culture*, (Rothblat, G.H. and Cristofalo, V.J. eds.), pp. 257–296, Academic Press, New York.
  - 51) Sata, R., Tsujii, H., Abe, H., Yamashita, S. and Hoshi, H. (1999): Fatty acid composition of bovine embryos cultured in serum-free and serum-containing medium during early embryonic development. *J. Reprod. Dev.*, 45, 97–103.
  - 52) Ferguson, E.M. and Leese, H.J. (1999): Triglyceride content of bovine oocytes and early embryos. *J. Reprod. Fertil.*, 116, 373–378.
  - 53) Kardymowicz, O. (1972): A method of vital staining for determining the viability of fertilized sheep ova stored in vitro. *Proc. 7th International Congress on Animal Reproduction & AI, Munich*, 1, 503–506.
  - 54) Schilling, E., Smidt, D., Sacher, B., Petac, D. and El Kaschab, S. (1979): Diagnosis of the viability of early bovine embryos by fluorescence microscopy. *Ann. Biol. Anim. Biochem. Biophys.*, 19, 1625–1629.
  - 55) Renard, J.P., Philippon, A. and Menezo, Y. (1980): In vitro uptake of glucose by bovine blastocysts. *J. Reprod. Fertil.*, 58, 161–164.
  - 56) Schilling, E., Niemann, H., Cheng, S.P. and Duepke, H.H. (1979): DAPI—a further fluorescence test for diagnosing the viability of early cow and rabbit embryos. *Zuchthyg.*



- 14, 170–172.
- 57) Elsdon, R.P., Nelson, L.D. and Seidel, G.E.J. (1978): Superovulating cows with follicle stimulating hormone and pregnant mare's serum gonadotrophin. *Theriogenology*, 9, 17–26.
  - 58) Lindner, G.M. and Wright, R.W.J. (1983): Bovine embryo morphology and evaluation. *Theriogenology*, 20, 407–416.
  - 59) Schneider, H.J.J., Castleberry, R.S. and Griffin, J.L. (1980): Commercial aspects of bovine embryo transfer. *Theriogenology*, 13, 73–85.
  - 60) Shea, B.F. (1981): Evaluating the bovine embryo. *Theriogenology*, 15, 13–42.
  - 61) Wright, J.M. (1981): Nonsurgical embryo transfer in cattle: embryo-recipient interactions. *Theriogenology*, 15, 43–56.
  - 62) Abe, H., Matsuzaki, S. and Hoshi, H. (2002): Ultrastructural differences in bovine morulae classified as high and low qualities by morphological evaluation. *Theriogenology*, 57, 1273–1283.
  - 63) Mills, R.M. and Brinster, R.L. (1967): Oxygen consumption of preimplantation mouse embryos. *Exp. Cell Res.*, 47, 337–344.
  - 64) Brinster, R.L. (1967): Carbon dioxide production from glucose by the preimplantation mouse embryo. *Exp. Cell Res.*, 47, 271–277.
  - 65) Wales, R.G., Whittingham, D.G., Hardy, K. and Craft, I.L. (1987): Metabolism of glucose by human embryos. *J. Reprod. Fertil.*, 79, 289–297.
  - 66) Rieger, D., Loskutoff, N.M. and Betteridge, K.J. (1992): Developmentally related changes in the uptake and metabolism of glucose, glutamine and pyruvate by cattle embryos produced in vitro. *Reprod. Fertil. Dev.*, 4, 547–557.
  - 67) Khurana, N.K. and Niemann, H. (2000): Energy metabolism in preimplantation bovine embryos derived in vitro or in vivo. *Biol. Reprod.*, 62, 847–856.
  - 68) Magnusson, C., Hillensjo, T., Hamberger, L. and Nilsson, L. (1986): Oxygen consumption by human oocytes and blastocysts grown in vitro. *Hum. Reprod.*, 1, 183–184.
  - 69) Magnusson, C., Hillensjo, T., Tsafiriri, A., Hultborn, R. and Ahren, K. (1977): Oxygen consumption of maturing rat oocytes. *Biol. Reprod.*, 17, 9–15.
  - 70) Nilsson, B., Magnusson, C., Widehn, S. and Hillensjo, T. (1982): Correlation between blastocyst oxygen consumption and trophoblast cytochrome oxidase reaction at initiation of implantation of delayed mouse blastocysts. *J. Embryol. Exp. Morphol.*, 71, 75–82.
  - 71) Houghton, F.D., Thompson, J.G., Kennedy, C.J. and Leese, H.J. (1996): Oxygen consumption and energy metabolism of the early mouse embryo. *Mol. Reprod. Dev.*, 44, 476–485.
  - 72) Thompson, J.G., Partridge, R.J., Houghton, F.D., Cox, C.I. and Leese, H.J. (1996): Oxygen uptake and carbohydrate metabolism by in vitro derived bovine embryos. *J. Reprod. Fertil.*, 106, 299–306.
  - 73) Donnay, I. and Leese, H.J. (1999): Embryo metabolism during the expansion of the bovine blastocyst. *Mol. Reprod. Dev.*, 53, 171–178.
  - 74) Overstrom, E.W., DUBY, R.T., Dobrinsky, J., Roche, J.F. and Boland, M.P. (1992): Viability and oxidative metabolism of the bovine blastocyst. *Theriogenology*, 37, 269.
  - 75) Smith, P.J.S., Hammar, K., Porterfield, D.M., Sanger, R.H. and Trimarchi, J.R. (1999): Self-referencing, non-invasive, ion selective electrode for single cell detection of transplasma membrane calcium flux. *Microsc. Res. Tech.*, 46, 398–417.
  - 76) Land, S.C., Porterfield, D.M., Sanger, R.H. and Smith, P.J.S. (1999): The self-referencing oxygen-selective microelectrode: detection of transmembrane oxygen flux from single cells. *J. Exp. Biol.*, 202, 211–218.
  - 77) Porterfield, D.M., Trimarchi, J.R., Keefe, D.L. and Smith, P.J.S. (1998): Characterization of oxygen and calcium fluxes from early mouse embryos and oocytes. *Biol. Bull.*, 195, 208–209.
  - 78) Trimarchi, J.R., Liu, L., Porterfield, D.M., Smith, P.J.S. and Keefe, D.L. (2000): Oxidative phosphorylation-dependent and -independent oxygen consumption by individual preimplantation mouse embryos. *Biol. Reprod.*, 62, 1866–1874.
  - 79) Shiku, H., Shiraishi, T., Ohya, H., Matsue, T., Abe, H., Hoshi, H. and Kobayashi, M. (2001): Oxygen consumption of single bovine embryos probed with scanning electrochemical microscopy. *Anal. Chem.*, 73, 3751–3758.